arXiv:1402.3096v1 [math.LO] 13 Feb 2014

Relations on FP-Soft Sets Applied to Decision Making Problems

Irfan Deli
Department of Mathematics, Faculty of Arts and Sciences
7 Aralık University, 79000 Kilis, Turkey
irfandeli@kilis.edu.tr

Naim Çağman
Department of Mathematics, Faculty of Arts and Sciences
Gaziosmanpaşa University, 60250 Tokat, Turkey
naim.cagman@gop.edu.tr

February 14, 2014

Abstract

In this work, we first define relations on the fuzzy parametrized soft sets and study their properties. We also give a decision making method based on these relations. In approximate reasoning, relations on the fuzzy parametrized soft sets have shown to be of a primordial importance. Finally, the method is successfully applied to a problems that contain uncertainties.

Keyword 0.1 *Soft sets, fuzzy sets, FP-soft sets, relations on FP-soft sets, decision making.*

1 Introduction

In 1999, the concept of soft sets was introduced by Molodtsov [25] to deal with problems that contain uncertainties. After Molodtsov, the operations of soft sets are given in [4, 23, 28] and studied their properties. Since then, based on these operations, soft set theory has developed in many directions and applied to wide variety of fields. For instance; on the theory of soft sets [2, 4, 5, 9, 20, 23, 24, 28], on the soft decision making [16, 17, 18, 21, 22, 27], on the fuzzy soft sets [7, 10, 11] and soft rough sets [16] are some of the selected works. Some authors have also studied the algebraic properties of soft sets, such as [1, 3, 6, 19, 26, 29, 30].

The FP-soft sets, firstly studied by Çağman *et al.* [8], is a fuzzy parameterized soft sets. Then, FP-soft sets theory and its applications studied in detail, for example [12, 13, 14]. In this paper, after given most of the fundamental

definitions of the operations of fuzzy sets, soft sets and FP-soft sets in next section, we define relations on FP-soft sets and we also give their properties in Section 3. In Section 4, we define symmetric, transitive and reflexive relations on the FP-soft sets. In Section 5, we construct a decision making method based on the FP-soft sets. We also give an application which shows that this methods successfully works. In the final section, some concluding comments are presented.

2 Preliminary

In this section, we give the basic definitions and results of soft set theory [25] and fuzzy set theory [31] that are useful for subsequent discussions.

Definition 2.1 *[31] Let U be the universe. A fuzzy set X over U is a set defined by a membership function μ_X representing a mapping*

$$\mu_X : U \to [0,1].$$

The value $\mu_X(x)$ for the fuzzy set X is called the membership value or the grade of membership of $x \in U$. The membership value represents the degree of x belonging to the fuzzy set X. Then a fuzzy set X on U can be represented as follows,

$$X = \{(\mu_X(x)/x) : x \in U, \mu_X(x) \in [0,1]\}.$$

Note that the set of all fuzzy sets on U will be denoted by $F(U)$.

Definition 2.2 *[15] t-norms are associative, monotonic and commutative two valued functions t that map from $[0,1] \times [0,1]$ into $[0,1]$. These properties are formulated with the following conditions:*

1. $t(0,0) = 0$ *and* $t(\mu_{X_1}(x), 1) = t(1, \mu_{X_1}(x)) = \mu_{X_1}(x),\ x \in E$
2. *If* $\mu_{X_1}(x) \leq \mu_{X_3}(x)$ *and* $\mu_{X_2}(x) \leq \mu_{X_4}(x)$, *then* $t(\mu_{X_1}(x), \mu_{X_2}(x)) \leq t(\mu_{X_3}x), \mu_{X_4}(x))$
3. $t(\mu_{X_1}(x), \mu_{X_2}(x)) = t(\mu_{X_2}(x), \mu_{X_1}(x))$
4. $t(\mu_{X_1}(x), t(\mu_{X_2}(x), \mu_{X_3}(x))) = t(t(\mu_{X_1}(x), \mu_{X_2})(x), \mu_{X_3}(x))$

Definition 2.3 *[15] t-conorms or s-norm are associative, monotonic and commutative two placed functions s which map from $[0,1] \times [0,1]$ into $[0,1]$. These properties are formulated with the following conditions:*

1. $s(1,1) = 1$ *and* $s(\mu_{X_1}(x), 0) = s(0, \mu_{X_1}(x)) = \mu_{X_1}(x), x \in E$
2. *if* $\mu_{X_1}(x) \leq \mu_{X_3}(x)$ *and* $\mu_{X_2}(x) \leq \mu_{X_4}(x)$, *then* $s(\mu_{X_1}(x), \mu_{X_2}(x)) \leq s(\mu_{X_3}(x), \mu_{X_4}(x))$
3. $s(\mu_{X_1}(x), \mu_{X_2}(x)) = s(\mu_{X_2}(x), \mu_{X_1}(x))$

4. $s(\mu_{X_1}(x), s(\mu_{X_2}(x), \mu_{X_3}(x))) = s(s(\mu_{X_1}(x), \mu_{X_2})(x), \mu_{X_3}(x))$

t-norm and t-conorm are related in a sense of lojical duality. Typical dual pairs of non parametrized t-norm and t-conorm are complied below:

1. Drastic product:

$$t_w(\mu_{X_1}(x), \mu_{X_2}(x)) = \begin{cases} min\{\mu_{X_1}(x), \mu_{X_2}(x)\}, & max\{\mu_{X_1}(x)\mu_{X_2}(x)\} = 1 \\ 0, & otherwise \end{cases}$$

2. Drastic sum:

$$s_w(\mu_{X_1}(x), \mu_{X_2}(x)) = \begin{cases} max\{\mu_{X_1}(x), \mu_{X_2}(x)\}, & min\{\mu_{X_1}(x)\mu_{X_2}(x)\} = 0 \\ 1, & otherwise \end{cases}$$

3. Bounded product:

$$t_1(\mu_{X_1}(x), \mu_{X_2}(x)) = max\{0, \mu_{X_1}(x) + \mu_{X_2}(x) - 1\}$$

4. Bounded sum:

$$s_1(\mu_{X_1}(x), \mu_{X_2}(x)) = min\{1, \mu_{X_1}(x) + \mu_{X_2}(x)\}$$

5. Einstein product:

$$t_{1.5}(\mu_{X_1}(x), \mu_{X_2}(x)) = \frac{\mu_{X_1}(x).\mu_{X_2}(x)}{2 - [\mu_{X_1}(x) + \mu_{X_2}(x) - \mu_{X_1}(x).\mu_{X_2}(x)]}$$

6. Einstein sum:

$$s_{1.5}(\mu_{X_1}(x), \mu_{X_2}(x)) = \frac{\mu_{X_1}(x) + \mu_{X_2}(x)}{1 + \mu_{X_1}(x).\mu_{X_2}(x)}$$

7. Algebraic product:

$$t_2(\mu_{X_1}(x), \mu_{X_2}(x)) = \mu_{X_1}(x).\mu_{X_2}(x)$$

8. Algebraic sum:

$$s_2(\mu_{X_1}(x), \mu_{X_2}(x)) = \mu_{X_1}(x) + \mu_{X_2}(x) - \mu_{X_1}(x).\mu_{X_2}(x)$$

9. Hamacher product:

$$t_{2.5}(\mu_{X_1}(x), \mu_{X_2}(x)) = \frac{\mu_{X_1}(x).\mu_{X_2}(x)}{\mu_{X_1}(x) + \mu_{X_2}(x) - \mu_{X_1}(x).\mu_{X_2}(x)}$$

10. Hamacher sum:

$$s_{2.5}(\mu_{X_1}(x), \mu_{X_2}(x)) = \frac{\mu_{X_1}(x) + \mu_{X_2}(x) - 2.\mu_{X_1}(x).\mu_{X_2}(x)}{1 - \mu_{X_1}(x).\mu_{X_2}(x)}$$

11. Minumum:

$$t_3(\mu_{X_1}(x), \mu_{X_2}(x)) = min\{\mu_{X_1}(x), \mu_{X_2}(x)\}$$

12. Maximum:

$$s_3(\mu_{X_1}(x), \mu_{X_2}(x)) = max\{\mu_{X_1}(x), \mu_{X_2}(x)\}$$

Definition 2.4 *[25]. Let U be an initial universe set and let E be a set of parameters. Then, a pair (F, E) is called a soft set over U if and only if F is a mapping or E into the set of aft subsets of the set U.*

In other words, the soft set is a parametrized family of subsets of the set U. Every set $F(\varepsilon)$, $\varepsilon \in E$, from this family may be considered as the set of ε-elements of the soft set (F, E), or as the set of ε-approximate elements of the soft set.

It is worth noting that the sets $F(\varepsilon)$ may be arbitrary. Some of them may be empty, some may have nonempty intersection.

In this definition, E is a set of parameters that are describe the elements of the universe U. To apply the soft set in decision making subset $A, B, C, ...$ of the parameters set E are needed. Therefore, Çağman and Enginoğlu [4] modified the definition of soft set as follows.

Definition 2.5 *[4] Let U be a universe, E be a set of parameters that are describe the elements of U, and $A \subseteq E$. Then, a soft set F_A over U is a set defined by a set valued function f_A representing a mapping*

$$f_A : E \to P(U) \text{ such that } f_A(x) = \emptyset \text{ if } x \in E - A \tag{1}$$

where f_A is called approximate function of the soft set F_A. In other words, the soft set is a parametrized family of subsets of the set U, and therefore it can be written a set of ordered pairs

$$F_A = \{(x, f_A(x)) : x \in E, f_A(x) = \emptyset \text{ if } x \in E - A\}$$

The subscript A in the f_A indicates that f_A is the approximate function of F_A. The value $f_A(x)$ is a set called *x-element* of the soft set for every $x \in E$.

Definition 2.6 *[8] Let F_X be a soft set over U with its approximate function f_X and X be a fuzzy set over E with its membership function μ_X. Then, a $FP-$soft sets Γ_X, is a fuzzy parameterized soft set over U, is defined by the set of ordered pairs*

$$\Gamma_X = \{(\mu_X(x)/x, f_X(x)) : x \in E\}$$

where $f_X : E \to P(U)$ such that $f_X(x) = \emptyset$ if $\mu_X(x) = 0$ is called approximate function and $\mu_X : E \to [0, 1]$ is called membership function of $FP-$soft set Γ_X. The value $\mu_X(x)$ is the degree of importance of the parameter x and depends on the decision-maker's requirements.

Note that the sets of all $FP-$soft sets over U will be denoted by $FPS(U)$.

3 Relations on the FP-Soft Sets

In this section, after given the cartesian products of two FP-soft sets, we define a relations on FP-soft sets and study their desired properties.

Definition 3.1 *Let $\Gamma_X, \Gamma_Y \in FPS(U)$. Then, a cartesian product of Γ_X and Γ_Y, denoted by $\Gamma_X \widehat{\times} \Gamma_Y$, is defined as*

$$\Gamma_X \widehat{\times} \Gamma_Y = \left\{ \left(\mu_{X \widehat{\times} Y}(x,y)/(x,y), f_{X \widehat{\times} Y}(x,y) \right) : (x,y) \in E \times E \right\}$$

where

$$f_{X \widehat{\times} Y}(x,y) = f_X(x) \cap f_Y(y)$$

and

$$\mu_{X \widehat{\times} Y}(x,y) = min\{\mu_X(x), \mu_Y(y)\}$$

Here $\mu_{X \widehat{\times} Y}(x,y)$ is a t-norm.

Example 3.2 *Let $U = \{u_1, u_2, u_3, u_4, u_5, u_6, u_7, u_8, u_9, u_{10}, u_{11}, u_{12}, u_{13}, u_{14}, u_{15}\}$, $E = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$, and $X = \{0.5/x_1, 0.7/x_2, 0.3/x_3, 0.9/x_4, 0.6/x_5\}$ and $Y = \{0.9/x_3, 0.1/x_6, 0.7/x_7, 0.3/x_8\}$ be two fuzzy subsets of E. Suppose that*

$$\begin{aligned}
\Gamma_X = \; & \Big\{ (0.5/x_1, \{u_1, u_3, u_4, u_6, u_7, u_8, u_{11}, u_{12}, u_{13}, u_{15}\}), (0.7/x_2, \{u_3, u_7, u_8, u_{14}, \\
& u_{15}\}), (0.3/x_3, \{u_1, u_2, u_4, u_5, u_6, u_9, u_{10}, u_{12}, u_{13}\}), (0.9/x_4, \{u_2, u_4, u_6, u_8, \\
& u_{12}, u_{13}\}), (0.6/x_5, \{u_3, u_4, u_6, u_7, u_9, u_{13}, u_{15}\}) \Big\} \\
\Gamma_Y = \; & \Big\{ (0.9/x_3, \{u_1, u_5, u_6, u_9, u_{10}, u_{13}\}), (0.1/x_6, \{u_3, u_5, u_7, u_8, u_9, u_{11}, u_{15}\}), \\
& (0.7/x_7, \{u_2, u_5, u_9, u_{10}, u_{11}, u_{14}\}), (0.3/x_8, \{u_2, u_5, u_8, u_{10}, u_{12}, u_{14}\}) \Big\}
\end{aligned}$$

Then, the cartesian product of Γ_X and Γ_Y is obtained as follows

$$\begin{aligned}
\Gamma_X \widehat{\times} \Gamma_Y = \; & \Big\{ (0.5/(x_1, x_3), \{u_1, u_6, u_{13}\}), (0.1/(x_1, x_6), \{u_3, u_7, u_8, u_{11}, u_{15}\}), \\
& (0.5/(x_1, x_7), \{u_{11}\}), (0.3/(x_1, x_8), \{u_8, u_{12}\}), (0.7/(x_2, x_3), \emptyset), \\
& (0.1/(x_2, x_6), \{u_3, u_7, u_8\}), (0.7(x_2, x_7), \{u_{14}\}), (0.3/(x_2, x_8), \\
& \{u_8, u_{14}\}), (0.3/(x_3, x_3), \{u_1, u_5, u_6, u_9, u_{10}, u_{13}\}), (0.1/(x_3, x_6), \\
& \{u_5, u_9\}), (0.3/(x_3, x_7), \{u_2, u_5, u_9, u_{10}\}), (0.3/(x_3, x_8), \\
& \{u_2, u_5, u_{10}, u_{12}\}), (0.9/(x_4, x_3), \{u_6\}), (0.1/(x_4, x_6), \emptyset), \\
& (0.7/(x_4, x_7), \{u_2, u_6\}), (0.3/(x_4, x_8), \{u_2, u_8, u_{12}\}), \\
& (0.6/(x_5, x_3), \{u_6, u_9, u_{13}\}), (0.1/(x_5, x_6), \{u_3, u_7, u_9, u_{11}, u_{15}\}), \\
& (0.6/(x_5, x_7), \{u_2\}), (0.3/(x_5, x_8), \emptyset) \Big\}
\end{aligned}$$

Definition 3.3 *Let $\Gamma_X, \Gamma_Y \in FPS(U)$. Then, an FP-soft relation from Γ_X to Γ_Y, denoted by R_F, is an FP-soft subset of $\Gamma_X \widehat{\times} \Gamma_Y$. Any FP-soft subset of $\Gamma_X \times \Gamma_Y$ is called a FP-relation on Γ_X.*

Note that if $\alpha = (\mu_X(x), f_X(x)) \in \Gamma_X$ and $\beta = (\mu_Y(y), f_Y(y)) \in \Gamma_Y$, then

$$\alpha R_F \beta \Leftrightarrow \left(\mu_{X\widehat{\times}Y}(x,y)/(x,y), f_{X\widehat{\times}Y}(x,y)\right) \in R_F$$

Example 3.4 *Let us consider the Example 3.2. Then, we define an FP-soft relation R_F, from Γ_Y to Γ_X, as follows*

$$\alpha R_F \beta \Leftrightarrow \mu_{X\widehat{\times}Y}(x_i, x_j)/(x_i, x_j)) \geq 0.3 \quad (1 \leq i, j \leq 3)$$

Then

$$\begin{aligned} R_F = \Big\{ & (0.5/(x_1, x_3), \{u_1, u_6, u_{13}\}), ((0.5/(x_1, x_7), \{u_{11}\}), (0.3/(x_1, x_8), \{u_8, \\ & u_{12}\}), (0.7(x_2, x_7), \{u_{14}\}), (0.3/(x_2, x_8), \{u_8, u_{14}\}), (0.3/(x_3, x_3), \{u_1, \\ & u_5, u_6, u_9, u_{10}, u_{13}\}), (0.3/(x_3, x_7), \{u_2, u_5, u_9, u_{10}\}), (0.3/(x_3, x_8), \{u_2, \\ & u_5, u_{10}, u_{12}\}), (0.9/(x_4, x_3), \{u_6\}), (0.7/(x_4, x_7), \{u_2, u_6\}), (0.3/(x_4, x_8), \\ & \{u_2, u_8, u_{12}\}), (0.6/(x_5, x_3), \{u_6, u_9, u_{13}\}), (0.6/(x_5, x_7), \{u_2\}) \Big\} \end{aligned}$$

Definition 3.5 *Let $\Gamma_X, \Gamma_Y \in FPS(U)$ and R_F be an FP-soft relation from Γ_X to Γ_Y. Then domain and range of R_F respectively is defined as*

$$\begin{aligned} D(R_F) &= \{\alpha \in F_A : \alpha R_F \beta\} \\ R(R_F) &= \{\beta \in F_B : \alpha R_F \beta\}. \end{aligned}$$

Example 3.6 *Let us consider the Example 3.4.*

$$\begin{aligned} D(R_F) = \Big\{ & (0.5/x_1, \{u_1, u_3, u_4, u_6, u_7, u_8, u_{11}, u_{12}, u_{13}, u_{15}\}), (0.7/x_2, \{u_3, u_7, \\ & u_8, u_{14}, u_{15}\}), (0.3/x_3, \{u_1, u_2, u_4, u_5, u_6, u_9, u_{10}, u_{12}, u_{13}\}), (0.9/x_4, \\ & \{u_2, u_4, u_6, u_8, u_{12}, u_{13}\}), (0.6/x_5, \{u_3, u_4, u_6, u_7, u_9, u_{13}, u_{15}\}) \Big\} \end{aligned}$$

$$\begin{aligned} R(R_F) = \Big\{ & (0.9/x_3, \{u_1, u_5, u_6, u_9, u_{10}, u_{13}\}), (0.7/x_7, \{u_2, u_5, \\ & u_9, u_{10}, u_{11}, u_{14}\}), (0.3/x_8, \{u_2, u_5, u_8, u_{10}, u_{12}, u_{14}\}) \Big\} \end{aligned}$$

Definition 3.7 *Let R_F be an FP-soft relation from Γ_X to Γ_Y. Then R_F^{-1} is from Γ_Y to Γ_X is defined as*

$$\alpha R_F^{-1} \beta = \beta R_F \alpha$$

Example 3.8 *Let us consider the Example 3.4. Then, ${R_F}^{-1}$ is from Γ_Y to Γ_X is obtained by*

$$\begin{aligned} {R_F}^{-1} = \Big\{ & (0.5/(x_3, x_1), \{u_1, u_6, u_{13}\}), ((0.5/(x_7, x_1), \{u_{11}\}), (0.3/(x_8, x_1), \{u_8, \\ & u_{12}\}), (0.7(x_7, x_2), \{u_{14}\}), (0.3/(x_8, x_2), \{u_8, u_{14}\}), (0.3/(x_3, x_3), \{u_1, \\ & u_5, u_6, u_9, u_{10}, u_{13}\}), (0.3/(x_7, x_3), \{u_2, u_5, u_9, u_{10}\}), (0.3/(x_8, x_3), \{u_2, \\ & u_5, u_{10}, u_{12}\}), (0.9/(x_3, x_4), \{u_6\}), (0.7/(x_7, x_4), \{u_2, u_6\}), (0.3/(x_8, x_4), \\ & \{u_2, u_8, u_{12}\}), (0.6/(x_3, x_5), \{u_6, u_9, u_{13}\}), (0.6/(x_7, x_5), \{u_2\}) \Big\} \end{aligned}$$

Proposition 3.9 *Let R_{F_1} and R_{F_2} be two FP-soft relations. Then*

1. $(R_{F_1}^{-1})^{-1} = R_{F_1}$
2. $R_{F_1} \subseteq R_{F_2} \Rightarrow R_{F_1}^{-1} \subseteq R_{F_2}^{-1}$

Proof:

1. $\alpha(R_{F_1}^{-1})^{-1}\beta = \beta R_{F_1}^{-1}\alpha = \alpha R_{F_1}\beta$
2. $\alpha R_{F_1}\beta \subseteq \alpha R_{F_2}\beta \Rightarrow \beta R_{F_1}^{-1}\alpha \subseteq \beta R_{F_2}^{-1}\alpha \Rightarrow R_{F_1}^{-1} \subseteq R_{F_2}^{-1}$

Definition 3.10 *If R_{F_1} is a fuzzy parametrized soft relation from Γ_X to Γ_Y and R_{F_2} is a fuzzy parametrized soft relation from Γ_Y to Γ_Z, then a composition of two FP-soft relations R_{F_1} and R_{F_2} is defined by*

$$\alpha(R_{F_1} \circ R_{F_2})\gamma = (\alpha R_{F_1}\beta) \wedge (\beta R_{F_2}\gamma)$$

Proposition 3.11 *Let R_{F_1} and R_{F_2} be two FP-soft relation from Γ_X to Γ_Y. Then, $(R_{F_1} \circ R_{F_2})^{-1} = R_{F_2}^{-1} \circ R_{F_1}^{-1}$*

Proof:

$$\begin{aligned}
\alpha(R_{F_1} \circ R_{F_2})^{-1})\gamma &= \gamma(R_{F_1} \circ R_{F_2})\alpha \\
&= (\gamma R_{F_1}\beta) \wedge (\beta R_{F_2}\alpha) \\
&= (\beta R_{F_2}\alpha) \wedge (\gamma R_{F_1}\beta) \\
&= (\alpha R_{F_2}^{-1}\beta) \wedge (\beta R_{F_1}^{-1}\gamma) \\
&= \alpha(R_{F_2}^{-1} \circ R_{F_1}^{-1})\gamma
\end{aligned}$$

Therefore we obtain
$(R_{F_1} \circ R_{F_2})^{-1} = R_{F_2}^{-1} \circ R_{F_1}^{-1}$

Definition 3.12 *An FP-soft relation R_F on Γ_X is said to be an FP-soft symmetric relation if $\alpha R_{F_1}\beta \Rightarrow \beta R_{F_1}\alpha, \forall \alpha, \beta \in \Gamma_X$.*

Definition 3.13 *An FP-soft relation R_F on Γ_X is said to be an FP-soft transitive relation if $R_F \circ R_F \subseteq R_F$, that is, $\alpha R_F \beta$ and $\beta R_F \gamma \Rightarrow \alpha R_F \gamma, \forall \alpha, \beta, \gamma \in \Gamma_X$.*

Definition 3.14 *An FP-soft relation R_F on Γ_X is said to be an FP-soft reflexive relation if $\alpha R_F \alpha, \forall \alpha \in \Gamma_X$.*

Definition 3.15 *An FP-soft relation R_F on Γ_X is said to be an FP-soft equivalence relation if it is symmetric, transitive and reflexive.*

Example 3.16 *Let $U = \{u_1, u_2, u_3, u_4, u_5, u_6, u_7, u_8\}$, $E = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ and $X = \{0.5/x_1, 0.7/x_2, 0.3/x_3\}$ be a fuzzy subsets over E. Suppose that*

$$\begin{aligned}
\Gamma_X = \Big\{ &(0.5/x_1, \{u_1, u_3, u_4, u_6, u_7, u_8\}), (0.7/x_2, \{u_3, u_7, u_8\}), \\
&(0.3/x_3, \{u_1, u_2, u_4, u_5, u_6, u_9\}) \Big\}
\end{aligned}$$

Then, a cartesian product on Γ_X is obtained as follows

$$\begin{aligned}\Gamma_X \widehat{\times} \Gamma_X = \quad & \Big\{(0.5/(x_1,x_1),\{u_1,u_3,u_4,u_6,u_7,u_8\}),\\ & (0.5/(x_1,x_2),\{u_3,u_7,u_8\}),(0.3/(x_1,x_3),\{u_1,u_4,u_6\}),\\ & (0.5/(x_2,x_1),\{u_3,u_7,u_8\}),(0.7/(x_2,x_2),\{u_3,u_7,u_8\}),\\ & (0.3/(x_3,x_1),\{u_1,u_4,u_6\}),(0.3/(x_3,x_3),\{u_1,u_2,u_4,u_5,u_6,u_9\})\Big\}\end{aligned}$$

Then, we get a fuzzy parametrized soft relation R_F on F_X as follows

$$\alpha R_F \beta \Leftrightarrow \mu_{X\widehat{\times}Y}(x_i,x_j)/(x_i,x_j)) \geq 0.3 \quad (1 \leq i,j \leq 3)$$

Then

$$\begin{aligned}R_F = \quad & \Big\{(0.5/(x_1,x_1),\{u_1,u_3,u_4,u_6,u_7,u_8\}),(0.5/(x_1,x_2),\{u_3,u_7,u_8\}),\\ & (0.3/(x_1,x_3),\{u_1,u_4,u_6\}),(0.5/(x_2,x_1),\{u_3,u_7,u_8\}),\\ & (0.7/(x_2,x_2),\{u_3,u_7,u_8\}),(0.3/(x_3,x_1),\{u_1,u_4,u_6\}),\\ & (0.3/(x_3,x_3),\{u_1,u_2,u_4,u_5,u_6,u_9\})\Big\}\end{aligned}$$

R_F on Γ_X is an FP-soft equivalence relation because it is symmetric, transitive and reflexive.

Proposition 3.17 *If R_F is symmetric if and only if R_F^{-1} is so.*

Proof: If R_F is symmetric, then $\alpha R_F^{-1}\beta = \beta R_F \alpha = \alpha R_F \beta = \beta R_F^{-1}\alpha$. So, R_F^{-1} is symmetric.

Conversely, if R_F^{-1} is symmetric, then $\alpha R_F \beta = \alpha (R_F^{-1})^{-1}\beta = \beta (R_F^{-1})\alpha = \alpha (R_F^{-1})\beta = \beta R_F \alpha$ So, R_F is symmetric.

Proposition 3.18 *R_F is symmetric if and only if $R_F^{-1} = R_F$*

Proof: If R_F is symmetric, then $\alpha R_F^{-1}\beta = \beta R_F \alpha = \alpha R_F \beta$. So, $R_F^{-1} = R_F$.

Conversely, if $R_F^{-1} = R_F$, then $\alpha R_F \beta = \alpha R_F^{-1}\beta = \beta R_F \alpha$. So, R_F is symmetric.

Proposition 3.19 *If R_{F_1} and R_{F_2} are symmetric relations on Γ_X, then $R_{F_1} \circ R_{F_2}$ is symmetric on Γ_X if and only if $R_{F_1} \circ R_{F_2} = R_{F_2} \circ R_{F_1}$*

Proof: If R_{F_1} and R_{F_2} are symmetric, then it implies $R_{F_1}^{-1} = R_{F_1}$ and $R_{F_2}^{-1} = R_{F_2}$. We have $(R_{F_1} \circ R_{F_2})^{-1} = R_{F_2}^{-1} \circ R_{F_1}^{-1}$. then $R_{F_1} \circ R_{F_2}$ is symmetric. It implies $R_{F_1} \circ R_{F_2} = (R_{F_1} \circ R_{F_2})^{-1} = R_{F_2}^{-1} \circ R_{F_1}^{-1} = R_{F_2} \circ R_{F_1}$.

Conversely, $(R_{F_1} \circ R_{F_2})^{-1} = R_{F_2}^{-1} \circ R_{F_1}^{-1} = R_{F_2} \circ R_{F_1} = R_{F_1} \circ R_{F_2}$. So, $R_{F_1} \circ R_{F_2}$ is symmetric.

Corollary 3.20 *If R_F is symmetric, then R_F^n is symmetric for all positive integer n, where $R_F^n = \underbrace{R_F \circ R_F \circ ... \circ R_F}_{n \text{ times}}$.*

Proposition 3.21 *If R_F is transitive, then R_F^{-1} is also transitive.*

Proof:

$$\begin{aligned}\alpha R_F^{-1}\beta &= \beta R_F\alpha \supseteq \beta(R_F \circ R_F)\alpha \\ &= (\beta R_F\gamma) \wedge (\gamma R_F\alpha) \\ &= (\gamma R_F\alpha) \wedge (\beta R_F\gamma) \\ &= (\alpha R_F^{-1}\gamma) \wedge (\gamma R_F^{-1}\beta) \\ &= \alpha(R_F^{-1} \circ R_F^{-1})\beta\end{aligned}$$

So, $R_F^{-1} \circ R_F^{-1} \subseteq R_F^{-1}$. The proof is completed.

Proposition 3.22 *If R_F is transitive then $R_F \circ R_F$ is so.*

Proof:

$$\begin{aligned}\alpha(R_F \circ R_F)\beta &= (\alpha R_F\gamma) \wedge (\gamma\text{-}R_F\beta) \\ &= \alpha(R_F \circ R_F)\gamma \wedge \gamma(R_F \circ R_F)\beta \\ &= \alpha(R_F \circ R_F \circ R_F \circ R_F)\beta\end{aligned}$$

So, $\alpha(R_F \circ R_F \circ R_F \circ R_F)\beta \subseteq \alpha(R_F \circ R_F)\beta$. The proof is completed.

Proposition 3.23 *If R_F is reflexive then R_F^{-1} is so.*

Proof: $\alpha R_F^{-1}\beta = \beta R_F\alpha \subseteq \alpha R_F\alpha = \alpha R_F^{-1}\alpha$ and $\beta R_F^{-1}\alpha = \alpha R_F\beta \subseteq \alpha R_F\alpha = \alpha R_F^{-1}\alpha$. The proof is completed.

Proposition 3.24 *If R_F is symmetric and transitive, then R_F is reflexive.*

Proof: Proof can be made easily by using Definition 4.1, Definition 4.2 and Definition 4.3.

Definition 3.25 *Let $\Gamma_X \in FPS(U)$, R_F be an FP-soft equivalence relation on Γ_X and $\alpha \in R_F$. Then, an equivalence class of α, denoted by $[\alpha]_{R_F}$, is defined as*

$$[\alpha]_{R_F} = \{\beta : \alpha R_F\beta\}.$$

Example 3.26 *Let us consider the Example 3.16. Then an equivalence class of*
$(x_1, \{u_1, u_3, u_4, u_6, u_7, u_8\})$ *will be as follows.*

$$[(0.5/x_1, \{u_1, u_3, u_4, u_6, u_7, u_8\})]_{R_F} = \left\{\begin{array}{l}(0.5/x_1, \{u_1, u_3, u_4, u_6, u_7, u_8\}), \\ (0.7/x_2, \{u_3, u_7, u_8\}), (0.3/x_3, \\ \{u_1, u_2, u_4, u_5, u_6, u_9\})\end{array}\right\}$$

4 Decision Making Method

In this section, we construct a soft fuzzification operator and a decision making method on FP-soft relations.

Definition 4.1 *Let $\Gamma_X \in FPS(U)$ and R_F be a FP-soft relation on Γ_X. Then fuzzification operator, denoted by s_{R_F}, is defined by*

$$s_{R_F} : R_F \to F(U), \quad s_{R_F}(X \times X, U) = \{\mu_{R_F}(u)/u : u \in U\}$$

where

$$\mu_{R_F}(u) = \frac{1}{|X \times X|} \sum_j \sum_i \mu_{R_F}(x_i, x_j)\chi(u)$$

and where

$$\chi(u) = \begin{cases} 1, & u \in f_{R_F}(x_i, x_j) \\ 0, & u \notin f_{R_F}(x_i, x_j) \end{cases}$$

Note that $|X \times X|$ is the cardinality of $X \times X$.

Now; we can construct a decision making method on FP-soft relation by the following algorithm;

1. construct a feasible fuzzy subset X over E,
2. construct a FP-soft set Γ_X over U,
3. construct a FP-soft relation R_F over Γ_X according to the requests,
4. calculate the fuzzification operator s_{R_F} over R_F,
5. select the objects, from s_{R_F}, which have the largest membership value.

Example 4.2 *A customer, Mr. X, comes to the auto gallery agent to buy a car which is over middle class. Assume that an auto gallery agent has a set of different types of car $U = \{u_1, u_2, u_3, u_4, u_5, u_6, u_7, u_8\}$, which may be characterized by a set of parameters $E = \{x_1, x_2, x_3, x_4\}$. For $i = 1, 2, 3, 4$ the parameters x_i stand for "safety", "cheap", "modern" and "large", respectively. If Mr. X has to consider own set of parameters, then we select a car on the basis of the set of customer parameters by using the algorithm as follows.*

1. *Mr X constructs a fuzzy sets X over E,*
   $X = \{0.5/x_1, 0.7/x_2, 0.3/x_3\}$
2. *Mr X constructs a FP-soft set Γ_X over U,*

$$\Gamma_X = \{(0.5/x_1, \{u_1, u_3, u_4, u_6, u_7, u_8\}), (0.7/x_2, \{u_3, u_7, u_8\}), (0.3/x_3, \{u_1, u_2, u_4, u_5, u_6, u_9\}$$

3. *the fuzzy parametrized soft relation R_F over Γ_X is calculated according to the Mr X's requests (The car must be a over middle class, it means the membership degrees are over 0.5),*

$$R_F = \Big\{(0.5/(x_1,x_1),\{u_1,u_3,u_4,u_6,u_7,u_8\}),(0.5/(x_1,x_2),\{u_3,u_7,u_8\}),(0.5/(x_2,x_1),\{u_3,u_7,u_8\}),(0.7/(x_2,x_2),\{u_3,u_7,u_8\})\Big\}$$

4. *the soft fuzzification operator s_{R_F} over R_F is calculated as follows*

$$s_{R_F} = \Big\{(0.055/u_1,0.0/u_2,0.244/u_3,0.055/u_4,0.0/u_5,0.055/u_6,0.244/u_7,0.244/u_8\}\Big\}$$

5. *now, select the optimum alternative objects u_3, u_7 and u_8 which have the biggest membership degree 0.244 among the others.*

5 Conclusion

We first gave most of the fundamental definitions of the operations of fuzzy sets, soft sets and FP-soft sets are presented. We then defined relations on FP-soft sets and studied some of their properties. We also defined symmetric, transitive and reflexive relations on the FP-soft sets. Finally, we construct a decision making method and gave an application which shows that this method successfully works. We have used a t-norm, which is minimum operator, the above relation. However, application areas the relations can be expanded using the above other norms in the future.

References

[1] Acar, U. Koyuncu, F. and Tanay, B. Soft sets and soft rings, Comput. Math. Appl. 59, 3458-3463, 2010.

[2] Ali, M.I. Feng, F. Liu, X. Min, W.K. and Shabir, M. On some new operations in soft set theory, Comput. Math. Appl. 57, 1547-1553, 2009.

[3] Aktaş, H. and Çağman, N. Soft sets and soft groups, Inform. Sci. 177, 2726-2735, 2007.

[4] Çağman, N. and Enginoğlu, S. Soft set theory and uni-int decision making, Eur. J. Oper. Res. 207, 848-855, 2010.

[5] Çağman, N. and Enginoğlu, S. Soft matrix theory and its decision making, Comput. Math. Appl. 59, 3308-3314, 2010.

[6] Çağman, N. Çıtak,, F. and Aktaş, H. Soft int-group and its applications to group theory, Neural Computing and Applications 21, 151-158, 2012.

[7] Çağman, N. and Enginoğlu, S. Fuzzy soft matrix theory and its applications in decision making, Iranian Journal of Fuzzy Systems, 9/1, 109-119, 2012.

[8] Çağman, N. Çıtak, F. and Enginoğlu, S. FP-soft set theory and its applications, Annals of Fuzzy Mathematics and Informatics 2/2, 219-226, 2011.

[9] Çağman, N. Karataş, S. and Enginoğlu, S. Soft Topology, Computers and Mathematics with Applications 62, 351 - 358, 2011.

[10] Çağman, N. Çıtak, F. and Enginoğlu, S. Fuzzy parameterized fuzzy soft set theory and its applications, Turkish Journal of Fuzzy Systems 1, 21-35, 2010.

[11] Çağman, N. Enginoğlu, S. and Çıtak, F. Fuzzy Soft Set Theory and Its Applications, Iran. J. Fuzzy Syst. 8/3, 137-147, 2011.

[12] Çağman, N. Deli, I. Means of FP-Soft Sets and its Applications, Hacettepe Journal of Mathematics and Statistics, (In Press)

[13] Çağman, N. Deli, I. Product of FP-Soft Sets and its Applications, Hacettepe Journal of Mathematics and Statistics, (In Press)

[14] Deli, I. Means of Fuzzy Parameterzed Soft Sets and its Applications (In Turkish), Masters Thesis, Gaziosmanpasa University, Graduate School of Natural and Applied Science, 2010.

[15] D. Dubois, and H. Prade, Fuzzy Set and Systems: Theory and Applications, Academic Press, New York, 1980.

[16] Feng, F. Liu, X.Y. Leoreanu-Fotea, V. Jun, Y.B. Soft sets and soft rough sets, Inform. Sci. 181/6, 1125-1137, 2011.

[17] Feng, F. Jun, Y.B. Liu, X. and Li, L. An adjustable approach to fuzzy soft set based decision making, J. Comput. App. Math. 234, 10-20, 2010.

[18] Feng, F. Li, L. Çağman, N. Generalized uni-int decision making schemes based on choice value soft sets, European Journal of Operational Research 220, 162-170, 2012.

[19] Jun, Y.B. Lee, K.J. and Khan, A. Soft ordered semigroups, Math. Logic Q. 56/1, 42-50, 2010.

[20] Karaaslan, F. Çağman N. and Enginoğlu, S. Soft Lattices, Journal of New Results in Science 1, 5-17, 2012.

[21] Kovkov, D.V. Kolbanov, V.M. and Molodtsov, D.A. Soft sets theory-based optimization, J. Comput. Sys. Sc. Int. 46/6, 872-880, 2007.

[22] Maji, P.K. Roy, A.R. and Biswas, R. An application of soft sets in a decision making problem, Comput. Math. Appl. 44, 1077-1083, 2002.

[23] Maji, P.K. Biswas, R. and Roy, A.R. Soft set theory, Comput. Math. Appl. 45, 555-562, 2003.

[24] Majumdar, P. and Samanta, S.K. Generalised fuzzy soft sets, Comput. Math. Appl. 59, 1425-1432, 2010.

[25] Molodtsov, D.A. Soft set theory-first results, Comput. Math. Appl. 37, 19-31, 1999.

[26] Park, C.H. Jun, Y.B. and Öztürk, M.A. Soft WS-algebras, Commun. Korean Math. Soc. 23/3, 313-324, 2008.

[27] Roy, A.R. and Maji, P.K. A fuzzy soft set theoretic approach to decision making problems, J. Comput. App. Math. 203, 412-418, 2007.

[28] Sezgin, A. and Atagün, A.O. On operations of soft sets, Comput. Math. Appl. 61/5, 1457-1467, 2011.

[29] Sezgin, A. Atagün, A.O. and Çağman N. Soft intersection near-rings with its applications, Neural Computing and Applications 21, 221-229, 2012.

[30] Sezgin, A. Atagün, A.O. and Çağman N. Union soft substructures of near-rings and N-groups, Neural Computing and Applications 21, 133-143, 2012.

[31] Zadeh, L.A. Fuzzy Sets, Inform. and Control 8, 338-353, 1965.

[32] Zimmermann, H.J. Fuzzy Set Theory and its applications. Kluwer Academic, 2th edition, Dordrecht, 1991.